

International Conference on **Obesity and Weight Loss**

November 6-8, 2017 Barcelona, Spain

Anthocyanin-rich Potato Improves Cardiovascular Risk Factors in Healthy Human Adults

C Tsang¹, Almoosawi S², Smail NF² and EAS Al-Dujaili²

¹Faculty of Health and Social Care, Edge Hill University, St Helens Road, Lancashire, UK

²Department of Dietetics, Nutrition and Biological Sciences, Queen Margaret University, Queen Margaret Drive, Edinburgh, UK

Arterial stiffness is an emerging risk factor for cardiovascular disease (CVD), and dietary polyphenols, particularly anthocyanins, may play an important role in mediating vascular tone. The present single-blind randomised cross-over placebo controlled study investigated the effect of consumption of an anthocyanin-rich potato, Purple Majesty (PM), in 14 healthy male and female adults. Participants consumed 200g PM containing 114 mg anthocyanins, or placebo (Osprey) with negligible anthocyanin content for 14 days, separated by a 1-week washout period. Non-invasive assessment of vascular tone (arterial stiffness) by pulse wave velocity (PWV) was determined in addition to systolic (SBP) and diastolic blood pressure (DBP), high density lipoproteins (HDL), low-density lipoproteins (LDL), triglycerides, glucose, insulin resistance (HOMA-IR) and c-reactive protein (CRP). Arterial stiffness was significantly reduced ($p=0.001$) following PM consumption. There were no significant changes with any other clinical parameter measured, and no changes were observed following placebo. PM contained higher levels of total phenolics, total anthocyanins and antioxidant capacity and daily consumption of PM over 14 days was well tolerated by participants. The findings from this study suggest improved vascular tone following daily consumption of PM and inclusion in the diet could provide a rich and valuable source of anthocyanins.