

## Impact of HCMV on HLA-I and Consequences on Gamma Delta T cell and Alpha Beta T cell Activation

**Layal Massara**

University of Bordeaux, College of Health Sciences, France

Cytomegalovirus (CMV), a Beta Herpes virus, is considered as a paradigm for viral evasion. It is an important opportunistic pathogen in immunocompromised patients and a major cause of congenital birth defects when acquired in utero. CMV encodes molecules to prevent antigen presentation to  $\alpha\beta$  T cells through inhibition of MHC Class I expression and to suppress NK cell functions by mimicking or down-regulating ligands of NK receptors (NKR). These evasion mechanisms are not expected to affect  $\gamma\delta$  T cells and, as a matter of fact, their response to CMV has been widely reported in many different physio-pathological contexts as well as in CMV-seropositive healthy donors (*Dechanet et al, 1999*)(*Scheper, 2013*).

Our aim was to understand how CMV induces  $\gamma\delta$  T cell response. We used recombinant adenoviruses expressing each of the four US genes, and a mutant HCMV deleted for these 4 genes (CMV-DUS). We observed an induction of HLA-I expression by the control adenovirus, and an inhibition by US2, US3 and US11. When using CMV-DUS, infected cells expressed much more native HLA-I than CMV-WT infected cells. Interestingly and in sharp contrast to  $\alpha\beta$  T cells,  $\gamma\delta$  T cells were activated to produce IFN $\gamma$  when cultured with fibroblasts infected with CMV-WT, but not when fibroblasts were infected with CMV-DUS. These results indicate that HLA-I molecules regulate  $\gamma\delta$  T cells through mechanisms that are under investigation in our team. The immune escape processes developed by CMV could thus promote  $\gamma\delta$  over  $\alpha\beta$  T cell response and explain the important response of  $\gamma\delta$  T cells to the virus in immune suppressed individuals.